

# HAGYARD CARE

## Equine Medical Institute

Hagyard-Davidson-McGee  
Associates, PLLC

Hagyard Field Care  
Davidson Surgery Center  
McGee Medicine Center  
McGee Fertility Center  
Hagyard Laboratory  
Hagyard Sport Horse

4250 Iron Works Pike  
Lexington, Kentucky 40511-8412  
info@hagyard.com  
www.hagyard.com

Phone (859) 255-8741  
Fax (859) 253-0196  
Medicine (859) 253-0002  
Surgery (859) 233-0026  
Laboratory (859) 259-3685

## YOUR HORSE HAS SALMONELLA BUT IS NOT SICK...

### SYMPTOMS

Your horse has been found to be passing ("shedding") salmonella bacteria in its manure, but does not have any associated clinical problems (eg. diarrhea). Shedding of salmonella bacteria is quite common even among healthy horses. Situations associated with stress (eg. transportation, hot weather, surgical procedures, exposure to antibiotics) make shedding more likely. Horses with a previous or recent history of intestinal disease (eg. colic, diarrhea) are at high risk of salmonella shedding. It is unlikely that your horse will experience any health problems related to this episode of salmonella shedding.

### REDUCING RISKS

Because salmonella can cause diarrhea and illness in horses, other animals, and human beings (especially children, senior citizens, or those with problems affecting the immune system), we suggest a few measures (below) to reduce the risk of infection to other animals or people in contact with your horse.

- ◆ Try not to subject your horse to stressful experiences. Hard exercise, long-distance, transportation, competition of any sort, and elective veterinary procedures should all be avoided for the next 3-4 weeks.
- ◆ Ensure that other horses do not come in contact with manure from this horse. This may mean keeping the horse in a separate paddock, cleaning its stall with separate or disinfected tools, and properly disposing of the manure.
- ◆ Wear disposable gloves when working with the horse. Carefully wash your hands when you are finished. Minimize contact between this horse and children.

During the 3-4 week period you may have your veterinarian culture the feces for salmonella to ensure the shedding has stopped. Up to 5 such cultures may be performed over a period of several days to confirm that the horse is not shedding. In the unlikely event that these cultures show that your horse is still shedding salmonella, please contact your veterinarian.

### SURVEILLANCE

*Hagyard Equine Medical Institute* has one of the highest equine caseloads in the world. Because we run a state-of-the-art hospital, we are dedicated to providing the best veterinary care possible to our patients. As part of our veterinary services, we run an infection control program to control and prevent infections in our patients during hospitalization. The main goal of this program is to develop an early warning system for detection of disease causing organisms and help prevent the potential spread of disease. This surveillance system entails performing bacterial cultures on samples from patients. A primary concern is for infectious organisms passed in manure (eg. salmonella, rotavirus and clostridium).

